arXiv:1212.1960v1 [hep-th] 10 Dec 2012

Electromagnetic Duality in Dyonic RN/CFT Correspondence

Bin Chen[1, 2, *] and Jia-ju Zhang[1, †]
[1] *Department of Physics, and State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, P.R. China*
[2] *Center for High Energy Physics, Peking University, Beijing 100871, P.R. China*

Four-dimensional dyonic Reissner-Nordström (RN) black hole is a solution of Einstein-Maxwell theory, which carries electromagnetic duality. It could be described holographically by two-dimensional conformal field theory (CFT) duals. By using the thermodynamics method, we identify two elementary CFT dual pictures, the electric one and the magnetic one, for the dyonic RN black hole. Moreover, we find that there are more general CFT dual pictures, generated by the $SL(2,Z)$ transformations acting on two elementary pictures. We show that this $SL(2,Z)$ symmetry is exactly the electromagnetic duality of the Einstein-Maxwell theory.

PACS: 04.70.Dy, 11.25.Hf

INTRODUCTION

One of central issues in quantum gravity is to understand microscopically the Bekenstein-Hawking entropy of a black hole. In string theory this has been achieved [1] for some kinds of supersymmetric Bogomol'nyi-Prasad-Sommerfield (BPS) black holes, whose entropy could be reproduced by counting the degeneracy of branes configuration. More precisely, the microscopical entropy is encoded in a two-dimensional (2D) conformal field theory (CFT), whose target spacetime is the moduli space of the branes configuration. Such kind of counting has been generalized to near-extremal and even non-extremal cases. In particular, for a class of five-dimensional (5D) and four-dimensional (4D) non-extreme multicharged black holes, their entropies could be written in an U-dual invariant form [2, 3]. Actually, even for the simplest case, there exist more than one dual pictures [4], the so-called long-string and short-string pictures, to describe the black hole. The existence of multiple holographic pictures indicates the underlying symmetry of the string theory.

In the past few years, the holographic 2D CFT descriptions of usual 4D rotating and charged black holes, which could not be embedded into string theory in a simple way, has been proposed and tested [5, 6]. In this so-called Kerr/CFT correspondence, people use completely different techniques, including the asymptotic symmetry group (ASG) analysis for the near horizon geometry of extremal black hole [5], and the hidden conformal symmetry in the low-frequency scattering off the non-extreme black hole [6], to investigate the holographic dual of the black hole. As these techniques have nothing to do with string theory or supersymmetry, they have been widely applied to many other kinds of black holes. Among them, the study on 4D Kerr-Newman and 5D Myers-Perry black holes is remarkable [7–11]. For each one of these two cases, there are multiple holographic pictures, which are generated by $SL(2,Z)$ transformations acting on two elementary ones. For example, for the 4D Kerr-Newman case, there are elementary J-picture based on the $U(1)$ rotation and elementary Q-picture based on $U(1)$ gauge symmetry [8, 10], from which the more generic pictures could be generated. This $SL(2,Z)$ group originates from the modular group of torus in the uplifted 5D metric [11].

Very recently, the Kerr/CFT correspondence was investigated from the point of view of thermodynamics [12–14]. From the first laws of thermodynamics of both the outer and inner horizons, one may compose the first laws of thermodynamics of a left-moving and a right-moving sectors. It turns out that these two sectors differs from the left-moving and right-moving sectors of dual CFT only by a scale factor. The scale factor could be determined from the first laws by considering the basic "quantized" unit of perturbation. In such a way, all the information of the dual CFT, including the temperatures and the central charges, could be read out easily. This method is remarkably effective, though intriguing. It not only has reproduced successfully the holographic pictures in well-established black hole/CFT correspondence [12], predicted the holographic picture of doubly-rotating black ring [13], but also helped to clarify the puzzles in RN/CFT correspondence [14]. Moreover, it allows us to read elementary CFT duals of the black hole in a simple way, and generate more generic dual pictures straightforwardly.

In this letter, we investigate the holographic pictures of dyonic RN black hole in 4D Einstein-Maxwell theory, using the thermodynamic method. This black hole carries both electronic and magnetic charges. Correspondingly, it has two elementary holographic pictures, the electric and magnetic pictures as we call. Using the first laws of thermodynamics, we may generate other holographic pictures by acting an $SL(2,Z)$ transformation on the elementary pictures. Unlike 4D Kerr-Newman or 5D Myers-Perry black holes, this $SL(2,Z)$ cannot be explained geometrically. Here we show that it originates from the electromagnetic duality of the Einstein-Maxwell theory.

DYONIC REISSNER-NORDSTRÖM BLACK HOLE

The dyonic RN black hole is a solution of the Einstein-Maxwell theory with the action $I = I_{EH} + I_{EM}$. I_{EH} is the Einstein-Hilbert action

$$I_{EH} = \frac{1}{16\pi G_4}\int d^4x\sqrt{-g}R, \tag{1}$$

and I_{EM} is the action of Maxwell theory in a curved background

$$I_{EM} = -\frac{1}{16\pi}\int d^4x\sqrt{-g}\left(F_{\mu\nu}F^{\mu\nu} + \frac{\theta e^2}{2\pi}F_{\mu\nu} * F^{\mu\nu}\right) \tag{2}$$

with $*$ being Hodge duality. We set $c = \hbar = 1$, $G_4 = \ell_p^2$ and use Gauss convention. To fully capture the features of electromagnetic duality, we have included the θ-term in the action. We may combine the two real coupling constants e, θ into a complex coupling parameter

$$\tau = \frac{\theta}{2\pi} + \frac{i}{e^2}. \tag{3}$$

The metric of the black hole is of the form

$$ds_4^2 = -N^2dt^2 + g_{rr}dr^2 + r^2(d\theta^2 + \sin^2\theta d\phi^2), \tag{4}$$

with

$$N^2 = \frac{1}{g_{rr}} = 1 - \frac{2G_4M}{r} + \frac{G_4Q^2}{r^2}. \tag{5}$$

Here M is the mass of the black hole, and

$$Q^2 = Q_e^2 + Q_m^2, \tag{6}$$

with $Q_{e,m}$ being the electric and magnetic charges respectively. As there is no globally well-defined gauge field for a dyon, we need to introduce two gauge fields to cover the whole spacetime manifold

$$A_{N,S} = -\frac{Q_e}{r}dt + Q_m(\cos\theta \mp 1)d\phi. \tag{7}$$

The upper minus sign applies to the north half-sphere and the lower plus sign applies to the south half-sphere.

The horizons locate at $r_\pm = G_4M \pm \sqrt{G_4^2M^2 - G_4Q^2}$. The entropy product of the outer and inner horizons is

$$\frac{S_+S_-}{4\pi^2} = \frac{Q^4}{4}, \tag{8}$$

which is mass-independent.

There is an electromagnetic duality in the action (2), which could be recast as

$$I_{EM} = -\frac{1}{32\pi}\int d^4x\sqrt{-g}\,\mathrm{Im}\mathcal{F}_{\mu\nu}\mathcal{G}^{\mu\nu}, \tag{9}$$

with $\mathcal{F}_{\mu\nu} = e(F_{\mu\nu} + i * F_{\mu\nu})$, $\mathcal{G}_{\mu\nu} = \tau\mathcal{F}_{\mu\nu}$. Under the $SL(2,Z)$ transformation

$$\begin{pmatrix} \mathcal{G}'_{\mu\nu} \\ \mathcal{F}'_{\mu\nu} \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}\begin{pmatrix} \mathcal{G}_{\mu\nu} \\ \mathcal{F}_{\mu\nu} \end{pmatrix}, \tag{10}$$

with $\alpha, \beta, \gamma, \delta \in Z$ and $\alpha\delta - \beta\gamma = 1$, the action is invariant. From $\mathcal{G}'_{\mu\nu} = \tau'\mathcal{F}'_{\mu\nu}$ it is easy to see that τ transform as

$$\tau' = \frac{\alpha\tau + \beta}{\gamma\tau + \delta}, \tag{11}$$

which is the feature of the electromagnetic duality. Correspondingly the charges transform as

$$\begin{pmatrix} e'Q_{e'} \\ \frac{Q_{m'}}{e'} - e'Q_{e'}\frac{\theta'}{2\pi} \end{pmatrix} = \begin{pmatrix} \delta & -\gamma \\ -\beta & \alpha \end{pmatrix}\begin{pmatrix} eQ_e \\ \frac{Q_m}{e} - eQ_e\frac{\theta}{2\pi} \end{pmatrix}. \tag{12}$$

Due to Witten's effect [15], the electric and the magnetic charges should be

$$\begin{aligned} Q_e &= N_ee - N_m\frac{e\theta}{2\pi}, \\ Q_m &= \frac{N_m}{e}, \end{aligned} \tag{13}$$

with $N_{e,m}$ being integers and transforming as

$$\begin{pmatrix} N_{e'} \\ N_{m'} \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}\begin{pmatrix} N_e \\ N_m \end{pmatrix}. \tag{14}$$

It can be verified easily that the quantity

$$Q^2 = Q_e^2 + Q_m^2 = \frac{1}{\mathrm{Im}\tau}|N_e - \tau N_m|^2 \tag{15}$$

is invariant under the above $SL(2,Z)$ transformation. Therefore the dyonic black hole geometry (4) does not change under the transformation of the electromagnetic duality. In other words, the geometry is the solution of different theories, which have different coupling constants and gauge potentials but are related to each other by the electromagnetic duality. This fact suggests that the dyonic black hole could be studied from the points of view of different theories, leading to different dual pictures.

DYONIC RN/CFT CORRESPONDENCE

For the dyonic RN black hole (4), it has the first laws of thermodynamics at the outer and the inner horizons

$$\begin{aligned} dM &= T_+dS_+ + \Omega_+^e dN_e + \Omega_+^m dN_m \\ &= -T_-dS_- + \Omega_-^e dN_e + \Omega_-^m dN_m. \end{aligned} \tag{16}$$

Here M is the mass of the black hole, $T_\pm$, $S_\pm$ are the temperatures and the entropies of the outer and the inner horizons. $N_{e,m}$ are two integer-valued charges related to $Q_{e,m}$ through (13), and $\Omega_{\pm}^{e,m}$ are their corresponding intensive quantities at the outer and the inner horizons. As clarified in [14], the quantization condition is essential to set up the holographic dual without ambiguity such that the integer-valued quantities N_e and N_m have been used in the above relations. It could be proved that the first law of the outer horizon always indicates that of the inner horizon, under reasonable assumptions [12], and if $T_+S_+ = T_-S_-$ is satisfied, then the entropy product S_+S_- is mass-independent. For convenience, one can define an entropy-product function

$$\mathcal{F} \equiv \frac{S_+S_-}{4\pi^2}, \tag{17}$$

with $\mathcal{F}$ being a function of the charges $N_{e,m}$ in our case.

After making the combinations[16–18]

$$\begin{aligned} T_{R,L} &= \frac{T_-T_+}{T_- \pm T_+}, \\ S_{R,L} &= \frac{1}{2}(S_+ \mp S_-), \\ \Omega_R^{e,m} &= \frac{T_-\Omega_+^{e,m} + T_+\Omega_-^{e,m}}{2(T_- + T_+)}, \\ \Omega_L^{e,m} &= \frac{T_-\Omega_+^{e,m} - T_+\Omega_-^{e,m}}{2(T_- - T_+)}, \end{aligned} \tag{18}$$

we can rewrite the first laws of the outer and inner horizons as the ones for the right- and left-moving sectors

$$\begin{aligned} \frac{1}{2}dM &= T_RdS_R + \Omega_R^e dN_e + \Omega_R^m dN_m \\ &= T_LdS_L + \Omega_L^e dN_e + \Omega_L^m dN_m. \end{aligned} \tag{19}$$

Keeping N_m invariant, i.e. considering the perturbation of the type $(dN_e, dN_m) = dN_e(1, 0)$, from the above first laws we could get

$$dN_e = \frac{T_L}{\Omega_R^e - \Omega_L^e}dS_L - \frac{T_R}{\Omega_R^e - \Omega_L^e}dS_R. \tag{20}$$

Now we can read the information of dual CFT. There are two remarkable facts on the above manipulations:

1. $S_{R,L}$ in (18) are exactly the entropies of the right- and left-moving sectors of CFT dual to the black hole[19, 20];
2. Keeping N_m invariant, we have the N_e picture, i.e. the electric picture. There is the first law of thermodynamics

   $$dN_e = T_L^e dS_L - T_R^e dS_R, \tag{21}$$

   with $T_{R,L}^e$ being exactly the right- and left-moving temperatures in the dual CFT[12, 14].

We assume that the CFT entropies could be reproduced by the Cardy formula

$$S_{R,L} = \frac{\pi^2}{3}c_{R,L}^e T_{R,L}^e, \tag{22}$$

from which we may derive the central charge. It could be shown easily that if $\mathcal{F}$ is mass-independent then the right- and left-moving sector central charges must be equal $c_R^e = c_L^e$. We substitute $S_\pm = S_L \pm S_R$ into (17) and take variations on both sides of the equation while keeping N_m invariant, then we get

$$\frac{\partial \mathcal{F}}{\partial N_e}dN_e = \frac{S_LdS_L - S_RdS_R}{2\pi^2}. \tag{23}$$

From the above equation, using the first law (21), the Cardy formula (22), and the fact that $c_R^e = c_L^e$, we find

$$c_{R,L}^e = 6\frac{\partial \mathcal{F}}{\partial N_e} = 6eQ_eQ^2. \tag{24}$$

These are the central charges of dual CFT in the electric picture [21]. Similarly, we could keep N_e invariant and obtain the N_m picture, i.e. the magnetic picture, in which the dual CFT is of the central charges

$$c_{R,L}^m = 6\frac{\partial \mathcal{F}}{\partial N_m} = 6Q^2\left(\frac{Q_m}{e} - eQ_e\frac{\theta}{2\pi}\right). \tag{25}$$

The relation between the entropy-product function $\mathcal{F}$ and the central charges is remarkable. For more general case, there is always the relation [22]

$$c_{R,L}^i = 6\frac{\partial \mathcal{F}}{\partial N_i}. \tag{26}$$

Here N_i is one of the integer-valued charges appearing in the first laws, and could be angular momentum, or other conserved $U(1)$ charges.

There is a simple way to understand the above elementary pictures. Actually the relation (19) tells us how the black hole responds to various kinds of the perturbations. If we just consider the perturbations carrying only the electric charges, or more precisely $(dN_e, dN_m) = dN_e(1, 0)$, the corresponding first law (21) gives us the temperatures $T_{R,L}^e$ and then the central charges $c_{R,L}^e$ in the electric picture. On the other hand, we may consider the perturbations carrying only magnetic charge, say $(dN_e, dN_m) = dN_m(0, 1)$, the first laws help us to read the magnetic picture, in which the CFT is of temperatures $T_{R,L}^m$ and central charges $c_{R,L}^m$. In general, we may consider the dyonic perturbations of the type $(dN_e, dN_m) = dN(a, b)$, with a, b being two coprime integers, then from the first laws (19) we could get

$$\begin{aligned} \frac{1}{2}dM &= T_RdS_R + \Omega_R^N dN \\ &= T_LdS_L + \Omega_L^N dN, \end{aligned} \tag{27}$$

with the intensive quantities conjugate to the integer-valued N being $\Omega^N_{R,L} = a\Omega^e_{R,L} + b\Omega^m_{R,L}$. Using the similar method, we can easily get the dyonic picture, or (a, b) picture, in which the CFT is of the central charges

$$c^{(a,b)}_{R,L} = ac^e_{R,L} + bc^m_{R,L}. \tag{28}$$

According to Bézout lemma, for every pair of coprime integers a, b there exists other pairs of coprime integers c, d so that $ad - bc = 1$. Thus the (a, b) picture CFT could be viewed as being generated from two elementary (1,0) and (0,1) pictures by a $SL(2, Z)$ transformation

$$\begin{pmatrix} c^{(a,b)}_{R,L} \\ c^{(c,d)}_{R,L} \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} c^{(1,0)}_{R,L} \\ c^{(0,1)}_{R,L} \end{pmatrix}, \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, Z).$$

In short, for the same dyonic RN black holes, there are many holographic descriptions, labeled by a pair coprime integers (a, b). These pictures are based on two elementary pictures and related to each other by $SL(2, Z)$ duality. Unlike the 4D Kerr-Newmann or 5D Myers-Perry black holes, this $SL(2, Z)$ could not be explained as the modular group of a two-torus. It has other interesting and profound origin.

$SL(2, Z)$ AS ELECTROMAGNETIC DUALITY

Using the electromagnetic duality there is another way of generating different CFT dual pictures. For a given dyonic black hole, we could describe it in different gravity theories, i.e. different Einstein-Maxwell theories, which are related by the electromagnetic duality. We begin with an Einstein-Maxwell theory describing the black hole with the electric and the magnetic charges given by (13), then we have the elementary (1,0) and (0,1) CFT pictures with the central charges (24), (25). Let us change the Einstein-Maxwell theory by an $SL(2, Z)$ duality described before

$$\tau' = \frac{\alpha\tau + \beta}{\gamma\tau + \delta}, \quad \begin{pmatrix} N_{e'} \\ N_{m'} \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} N_e \\ N_m \end{pmatrix}. \tag{29}$$

Then we get a gravity theory whose (1,0) picture central charges are

$$c^{e'}_{R,L} = 6e'Q_{e'}Q^2. \tag{30}$$

Note that there is no prime in Q^2 because it is $SL(2, Z)$ invariant. From the transformation (12), we get the central charges

$$c^{e'}_{R,L} = 6Q^2 \left[\delta e Q_e - \gamma \left(\frac{Q_m}{e} - eQ_e \frac{\theta}{2\pi} \right) \right]. \tag{31}$$

Comparing the results with (28), we find that we could get the same CFT picture by setting $\delta = a, \gamma = -b$.

We stress that (28) and (31) were obtained independently from two apparently different ways. The former was got by the $SL(2, Z)$ duality on the CFT theories, while the latter was got from the electromagnetic duality of the gravity theories. However it turns out the $SL(2, Z)$ duality in (28) originates from the electromagnetic duality in the gravity theory. Let us consider the dyonic (a, b)-type perturbation with $(dN_e, dN_m) = dN(a, b)$ in the original theory. After the $SL(2, Z)$ transformations (29) with

$$\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} d & -c \\ -b & a \end{pmatrix} \tag{32}$$

we have $(dN_{e'}, dN_{m'}) = dN(1, 0)$ which is just a (1,0) type perturbation in the "new" theory. Consequently from "new" theory point of view we have a (1,0) picture of the central charges from (31), in exact match with (28).

Furthermore, the electromagnetic duality in the gravity theory allows us to read the central charges by uplifting the solution to five dimension. For the gauge potential with both electric and magnetic charges, it is not clear how to uplift the metric. However, using the duality, it is always possible to consider the configuration in a theory with only pure electric field. In this case, the metric and gauge potential may be uplifted to a 5D metric, from which the central charges could be computed [14].

CONCLUSION AND DISCUSSION

In this letter we investigated the CFT duals of the four-dimensional RN black hole. It turned out that the electromagnetic duality played a subtle but important role in setting up generic holographic CFT duals of the black hole. It allowed us to study the same black hole background from different gravity theories, and find different pictures related to each other by the duality.

In four-dimensional Einstein-Maxwell gravity theory, there is dyonic Kerr-Newman black hole, which carries not only electric and magnetic charges but also angular momentum. From thermodynamics method, there seem to be three elementary pictures, from which the general dual pictures could be generated through $SL(3, Z)$ transformations. This $SL(3, Z)$ group could be understood as a kind of U-duality, combining the modular group of two-torus and the electromagnetic duality group. It would be interesting to understand this duality group more clearly.

The notion of electromagnetic duality is new in the context of 4D Einstein-Maxwell theory, but is well-known in string theory [23, 24]. The study in this paper indicates that the duality in the holographic descriptions of the black holes could be related to various duality in string theory. It has been found in [2, 3] that there are classes of black holes whose entropies could be written as U-dual invariant forms. This suggests that these black holes should have multi-fold holographic duals. It would be interesting to understand these duals more clearly from thermodynamics points of view [22]. On the other hand, even though the thermodynamics method is quite effective and robust, it deserves more investigation from microscopical point of view. We expect that string theory may shed light on this issue.

Acknowledgments

The work was in part supported by NSFC Grant No. 10975005, 11275010. JJZ was also in part supported by Scholarship Award for Excellent Doctoral Student granted by Ministry of Education of China.

---

[*] Email: bchen01@pku.edu.cn
[†] Email: jjzhang@pku.edu.cn

[1] A. Strominger and C. Vafa, Phys. Lett. B **379**, 99 (1996).
[2] G. T. Horowitz, J. M. Maldacena and A. Strominger, Phys. Lett. B **383**, 151 (1996).
[3] G. T. Horowitz, D. A. Lowe and J. M. Maldacena, Phys. Rev. Lett. **77**, 430 (1996).
[4] J. M. Maldacena and L. Susskind, Nucl. Phys. B **475**, 679 (1996).
[5] M. Guica, T. Hartman, W. Song and A. Strominger, Phys. Rev. D **80**, 124008 (2009).
[6] A. Castro, A. Maloney and A. Strominger, Phys. Rev. D **82**, 024008 (2010).
[7] H. Lu, J. Mei and C. N. Pope, JHEP **0904**, 054 (2009).
[8] T. Hartman, K. Murata, T. Nishioka and A. Strominger, JHEP **0904**, 019 (2009).
[9] C. Krishnan, JHEP **1007**, 039 (2010).
[10] C. M. Chen, Y. M. Huang, J. R. Sun, M. F. Wu and S. J. Zou, Phys. Rev. D **82**, 066004 (2010).
[11] B. Chen and J. -j. Zhang, Nucl. Phys. B **856**, 449 (2012); JHEP **1108**, 114 (2011).
[12] B. Chen, S. -x. Liu, and J. -j. Zhang, JHEP **1211**, 017 (2012).
[13] B. Chen and J. -j. Zhang, JHEP **1211**, 022 (2012).
[14] B. Chen and J. -j. Zhang, to appear.
[15] E. Witten, Phys. Lett. B **86**, 283 (1979).
[16] M. Cvetic and F. Larsen, Phys. Rev. D **56**, 4994 (1997).
[17] M. Cvetic and F. Larsen, Nucl. Phys. B **506**, 107 (1997).
[18] M. Cvetic and F. Larsen, JHEP **0909**, 088 (2009).
[19] M. Cvetic and D. Youm, Phys. Rev. D **54**, 2612 (1996).
[20] F. Larsen, Phys. Rev. D **56**, 1005 (1997).
[21] C. -M. Chen, Y. -M. Huang, J. -R. Sun, M. -F. Wu and S. -J. Zou, Phys. Rev. D **82**, 066003 (2010).
[22] B. Chen and J. -j. Zhang, work in progress.
[23] M. Cvetic and D. Youm, Nucl. Phys. Proc. Suppl. **46**, 56 (1996).
[24] M. Cvetic and A. A. Tseytlin, Phys. Lett. B **366**, 95 (1996).